

Symptoms and Medication Use in Children with Asthma and Traffic-Related Sources of Fine Particle Pollution

Janneane F. Gent,¹ Petros Koutrakis,² Kathleen Belanger,¹ Elizabeth Triche,¹ Theodore R. Holford,¹ Michael B. Bracken,¹ and Brian P. Leaderer¹

¹Center for Perinatal, Pediatric and Environmental Epidemiology, Department of Epidemiology and Public Health, Yale University School of Medicine, New Haven, Connecticut, USA; ²Harvard University School of Public Health, Boston, Massachusetts, USA

BACKGROUND: Exposure to ambient fine particles [particulate matter ≤ 2.5 μm diameter (PM_{2.5})] is a potential factor in the exacerbation of asthma. National air quality particle standards consider total mass, not composition or sources, and may not protect against health impacts related to specific components.

OBJECTIVE: We examined associations between daily exposure to fine particle components and sources, and symptoms and medication use in children with asthma.

METHODS: Children with asthma ($n = 149$) 4–12 years of age were enrolled in a year-long study. We analyzed particle samples for trace elements (X-ray fluorescence) and elemental carbon (light reflectance). Using factor analysis/source apportionment, we identified particle sources (e.g., motor vehicle emissions) and quantified daily contributions. Symptoms and medication use were recorded on study diaries. Repeated measures logistic regression models examined associations between health outcomes and particle exposures as elemental concentrations and source contributions.

RESULTS: More than half of mean PM_{2.5} was attributed to traffic-related sources motor vehicles (42%) and road dust (12%). Increased likelihood of symptoms and inhaler use was largest for 3-day averaged exposures to traffic-related sources or their elemental constituents and ranged from a 10% increased likelihood of wheeze for each 5- $\mu\text{g}/\text{m}^3$ increase in particles from motor vehicles to a 28% increased likelihood of shortness of breath for increases in road dust. Neither the other sources identified nor PM_{2.5} alone was associated with increased health outcome risks.

CONCLUSIONS: Linking respiratory health effects to specific particle pollution composition or sources is critical to efforts to protect public health. We associated increased risk of symptoms and inhaler use in children with asthma with exposure to traffic-related fine particles.

KEY WORDS: childhood asthma, fine particle pollution, PM_{2.5}, respiratory morbidity, source apportionment, traffic pollution. *Environ Health Perspect* 117:1168–1174 (2009). doi:10.1289/ehp.0800335 available via <http://dx.doi.org/> [Online 31 March 2009]

Fine particles [particles ≤ 2.5 μm diameter (PM_{2.5})] originate from both local and regional pollution sources. A large fraction of PM_{2.5} mass is associated with the combustion of fuels such as gasoline, diesel, coal, oil, and wood. PM_{2.5} mass also includes crustal material (road dust and soil particles), sea salt, and biological particles. Factor analysis and source apportionment techniques [Brown et al. 2007; U.S. Environmental Protection Agency (EPA) 2008; Watson et al. 2002] have been used for many years to help policy regulators identify sources of particle pollution. These techniques have been used to investigate associations between sources of particle pollution and health effects including studies of adult mortality (Laden et al. 2000; Mar et al. 2000; Ozkaynak and Thurston 1987; Taino et al. 2005), adult emergency department visits (Sarnat et al. 2008), and adult and pediatric hospital admissions (Andersen et al. 2007). Recently, one study associated specific sources of fine particles with respiratory morbidity in adults with asthma (Penttinen et al. 2006).

PM_{2.5} mass concentration has been associated with adverse health effects in children, particularly children with asthma: increased respiratory morbidity and/or medication use (Gent et al. 2003; Johnston et al. 2006; Mar

et al. 2004; Rabinovitch et al. 2006; Ranzi et al. 2004; Rodriguez et al. 2007; Slaughter et al. 2003); increased number of doctor visits for respiratory illness (Chardon et al. 2007; Hertz-Picciotto et al. 2007; Pino et al. 2004); and decreased lung function (Delfino et al. 2004; Hong et al. 2007; Lewis et al. 2005; Moshhammer et al. 2006; Trenga et al. 2006). Linking respiratory health effects to specific particle pollution sources is critical to efforts to develop cost-effective strategies to protect public health, particularly the respiratory health of vulnerable members of the population such as children with asthma.

Annual PM_{2.5} levels at a monitoring site in New Haven, Connecticut, where subjects in the present study live, exceed the U.S. EPA annual standard of 15 $\mu\text{g}/\text{m}^3$ (U.S. EPA 2009). We examined associations between daily exposure to PM_{2.5} components and sources identified through factor analysis and source apportionment, and daily symptoms and medication use in children with asthma. Some of the results of this study have been previously reported in the form of an abstract (Gent et al. 2008).

Methods

Subjects. Subjects were drawn from a larger study population 4–12 years of age when

enrolled between 2000 and 2003. Eligibility criteria were physician-diagnosed asthma and symptoms or medication use in the previous 12 months. Subjects in the present analysis ($n = 149$) were restricted to those residing in New Haven County within 30 km of the air quality monitor. Study participants included siblings with asthma (one per family; $n = 88$, 59%) of infants enrolled in a birth cohort to study the development of asthma (Belanger et al. 2003), patients recruited from the Yale Pediatric Asthma Clinic ($n = 43$, 29%), and children with asthma identified through a survey on childhood asthma conducted in the New Haven public schools ($n = 18$, 12%). In addition to abundant mobile sources of air pollution (e.g., interstate highways 95, 91, 84), New Haven is home to stationary sources including small manufacturing enterprises, power plants, and a harbor handling ships that deliver heating oil for New England (U.S. EPA 2008). New Haven County, population 824,008, covers an area of 2,233 km² (863 mi²) (U.S. Census Bureau 2000). Yale University Human Investigation Committee approved the study, and all respondents (mothers of study subjects) gave informed consent prior to participation.

Data collection. During the enrollment visit, a research assistant administered a questionnaire to the child's mother to collect demographic, medical history, and home environment information. Mothers recorded daily symptoms [wheeze, persistent cough (defined as coughing throughout the day), shortness of breath, chest tightness, night symptoms] and medication use (rescue medication including bronchodilators, and maintenance medication including steroids, cromolyn sodium, and leukotriene inhibitors) on study diaries and reported this information during monthly telephone interviews. Dates away from New Haven during the follow-up

Address correspondence to J.F. Gent, Yale Center for Perinatal, Pediatric and Environmental Epidemiology, One Church Street, 6th Floor, New Haven, CT 06510 USA. Telephone: (203) 764-9375. Fax: (203) 764-9378. E-mail: janneane.gent@yale.edu

Supplemental Material is available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>. This study was funded by National Institutes of Health grants ES011013 and ES05410.

The authors declare they have no competing financial interests.

Received 29 October 2008; accepted 31 March 2009.

period were reported at an exit interview and were not used in the analysis.

Exposure measurements. Filter analysis. PM_{2.5} filter samples ($n = 1,181$) were collected between 1 August 2000 and 3 February 2004 from the Stiles Street monitoring site (98% of the filters) or the State Street site (within 3 km). Filters were analyzed by X-ray fluorescence (XRF) (Desert Research Institute, Reno, Nevada) (Watson et al. 1999). A list of the 51 elements analyzed is presented in the Supplemental Material, Table 1 (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>).

Light reflectance analysis. Before XRF, elemental carbon (EC) concentrations were determined using an optical reflectance technique (Cyrys et al. 2003; Janssen et al. 2001; Kinney et al. 2000). See Supplemental Material for further details (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>).

Source apportionment. We analyzed daily concentrations of 17 elements [see Supplemental Material, Table 1 (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>)] as well as EC using principal component analysis with an orthogonal (varimax) rotation in SAS (version 9.1; SAS Institute Inc., Cary, NC) (Kavouras et al. 2001; Thurston and Spengler 1985). We calculated daily scores for each factor by multiplying the normalized elemental concentrations by the respective standardized elemental scoring coefficient and factor, then summing across all component products. Scores were rescaled to concentrations in micrograms per cubic meter by regressing total PM_{2.5} concentration on factor scores and obtaining the product of each factor score with its regression coefficient. Rescaled factor scores, or daily source contributions, were used as exposures in the health effects analyses.

Six sources of PM_{2.5} were identified based on loadings (or correlations) of elements on each of the six factors determined by factor analysis. The factor with high loadings for EC and trace elements zinc, lead, and copper was attributed to motor vehicle emissions. The factor road dust was associated with elements of terrestrial origin such as silicon, iron, aluminum, and calcium as well as barium, which is a tracer for tire wear particles. The sulfur factor represents particles from regional sources and was highly correlated with the elements sulfur and phosphorus. The factor with a high potassium loading was attributed to biomass burning. The oil factor represents oil combustion emissions from power plants and home heating and was characterized by the presence of vanadium and nickel. Finally, the sea salt factor was associated with the elements sodium and chlorine.

Data analysis. The association between daily elemental concentrations or source

contributions and daily respiratory symptoms and rescue medication use was examined with repeated measures logistic regression (SAS Institute Inc.). Models used generalized estimating equations (GEE) specifying a 1-day lag autoregressive structure (AR1) for the health outcomes correlation matrices. All models were adjusted for season using sine and cosine functions of date to capture annual cycles [i.e., $\cos((\text{date}/365) \times 2\pi)$ and $\sin((\text{date}/365) \times 2\pi)$], day of week (for weekly cycles), and date (to control for time trends over the 4-year course of the study). Because sources were uncorrelated by design (orthogonal rotation of factors), all six sources were included simultaneously in health effects models. Gaseous co-pollutants (nitrogen dioxide, carbon monoxide, sulfur dioxide, ozone) were examined separately in models that also included all six sources. Models included year-round data except for the co-pollutant model with O₃, which is monitored only in warm months. Health effects of elements included in the factor analysis, all six sources, and PM_{2.5} were examined for same-day exposures [lag0 (L0)] as well as exposures lagged by 1 or 2 days (L1, L2) or the mean of lagged days 0–2 (L02). Tests for goodness of fit were performed using the Hosmer–Lemeshow statistic for logistic regression.

Results

Demographic characteristics of study participants are shown in Table 1. Most subjects were male (60%), ≥ 8 years of age (53%) (mean \pm SD = 8 ± 2 years), and self-reported as white (56%). More than one-third of the adult respondents had a college degree (36%). Environmental tobacco smoke exposure was reported in 6% of the homes. Subjects participated for a mean of 313 ± 92 days [median 366; interquartile range

(IQR) 54 days]. The distribution of daily symptoms and medication use is shown in Table 2. The most frequently experienced symptom was persistent cough (reported by 86%), followed by wheeze (76%), which was experienced by subjects for a median of 19 days (IQR 29) or 8 days (IQR 24), respectively, of the year of follow-up. Subjects experienced shortness of breath or chest tightness (reported by two-thirds of the subjects) less frequently for a median of 4 (IQR 14) days each, and used short-acting inhalers (reported by 80%) for a median of 18 (IQR 47) days. Nearly one-third of the subjects had moderate to severe persistent asthma during the year-long follow-up study (Table 2). All subjects resided 0.9–27 km from the air

Table 2. Distribution of daily symptoms, daily inhaler use, and mean asthma severity during a year-long prospective study ($n = 149$), New Haven, Connecticut, August 2000–January 2004.

Variable	No. (%)
Symptoms and inhaler use during 12-month study period (days)	
Wheeze	
None	35 (23.5)
1–7	36 (24.2)
8–14	22 (14.8)
15–21	15 (10.1)
22–29	11 (7.4)
≥ 30	30 (20.1)
Persistent cough	
None	20 (13.5)
1–7	31 (21.0)
8–14	12 (8.1)
15–21	17 (11.5)
22–29	21 (14.2)
≥ 30	47 (31.8)
Shortness of breath	
None	46 (30.9)
1–7	43 (28.9)
8–14	23 (15.4)
15–21	9 (6.0)
22–29	9 (6.0)
≥ 30	19 (12.8)
Chest tightness	
None	49 (32.9)
1–7	44 (29.5)
8–14	20 (13.4)
15–21	9 (6.0)
22–29	7 (4.7)
≥ 30	20 (13.4)
Short-acting inhaler use	
None	29 (19.5)
1–7	28 (18.8)
8–14	14 (9.4)
15–21	11 (7.4)
22–29	11 (7.4)
≥ 30	56 (37.6)
Asthma severity (GINA score) ^a	
0 No symptoms/medication	5 (3.4)
1 Intermittent	66 (44.9)
2 Mild persistent	29 (19.7)
3 Moderate persistent	30 (20.4)
4 Severe persistent	17 (11.6)

^aA five-point asthma severity score [from 0 (no symptoms or medication use) to 4 (severe persistent)], based on the Global Initiative for Asthma guidelines (U.S. Department of Health and Human Services 2002) was calculated for each month of the study. The mean severity was the arithmetic mean of the 12 monthly severity scores.

Table 1. Personal characteristics for children participating in a year-long prospective study ($n = 149$), New Haven, Connecticut, August 2000–January 2004.

Enrollment characteristic	No. (%)
Sex	
Male	89 (59.7)
Female	60 (40.3)
Ethnicity	
White	84 (56.4)
Black	36 (24.2)
Hispanic	27 (18.1)
Other	2 (1.3)
Age at enrollment (years)	
< 8	70 (47.0)
≥ 8	79 (53.0)
Mother's education (years) ^a	
< 12	13 (8.7)
12–15	83 (55.7)
≥ 15	53 (36.0)
Environmental exposure to tobacco smoke	
No	140 (94.0)
Yes	9 (6.0)

^aThe adult respondent was the subject's mother in all but two homes.

pollution monitor site (mean \pm SD = 10.2 \pm 8.4 km). The mean number of subjects under observation each day was 36 \pm 8 and ranged from 10 to 49. The mean number of subjects experiencing symptoms or using short-acting inhalers each day was as follows: shortness of breath or chest tightness 3 \pm 3%; wheeze 5 \pm 4%; persistent cough 8 \pm 5%; and inhaler use 12 \pm 6% (Figure 1A,B).

Mean level of PM_{2.5} during the study was 17.0 \pm 9.8 $\mu\text{g}/\text{m}^3$, which is above the U.S. annual standard for fine particles of 15 $\mu\text{g}/\text{m}^3$ (U.S. EPA 2009). The contributions from each

source to PM_{2.5} mass and to elemental components of PM_{2.5} are given in Table 3. The mean estimated and measured mass are given as well as the percent root mean square error (%RMSE), defined as $100 \times (|\text{estimated concentration} - \text{measured concentration}| / \text{measured concentration})$. Elemental components of PM_{2.5} are grouped according to the source with which they were most strongly associated. For example, the motor vehicle source contributed 62% of the estimated mean EC mass (961.3 of 1554.8 ng/m³) and 77% of the mean Zn mass (22.3 of 29.0 ng/m³). On average, over

the 4 years of the study, over half of daily PM_{2.5} came from the two traffic-related sources motor vehicles [median (IQR) 42.5% (26.7%)] and road dust [12.3% (13.8%)] (bottom of Table 2). An additional 28.7% (30.3%) of daily PM_{2.5} came from regional sulfur sources.

The associations between PM_{2.5} elemental components and daily symptoms or medication use are presented in Table 4. Individual elements in fine particle mass may come from more than one source but are listed in Table 4 according to the source with which they were most strongly associated (Table 3). Associations for same-day exposure (L0) and exposures averaged over the same day and previous 2 days (L02) are shown in Table 4. Exposures lagged by 1 or 2 days (L1, L2) are included in Table 2 in the Supplemental Material (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>). In general, trace elements originating from the motor vehicle, road dust, biomass burning, and oil sources are associated with symptoms and/or medication use. For example, a significantly increased likelihood of wheeze, shortness of breath, chest tightness, or short-acting inhaler use is associated with each 1,000-ng/m³ increase in EC. The strongest associations were found for the 3-day averaged exposures to the elemental constituents of road dust. No associations were found between individual elements associated with sulfur or sea salt sources and symptoms or medication use. Total PM_{2.5} was not significantly associated with any symptoms or medication use [see Supplemental Material, Table 3 (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>)].

Daily source contributions to PM_{2.5} are illustrated for the 4 years of the study in Figure 1C–H. Results of logistic regression models examining associations between health outcomes and PM_{2.5} sources are shown in Table 5 for same-day exposures (Table 5, L0 model) and exposures averaged over the same day and previous 2 days (Table 5, L02 model). Source exposures lagged by 1 or 2 days (L1, L2) are included in the Supplemental Material, Table 3 (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>). For same-day or 3-day averaged exposures, each 5- $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} mass concentration coming from either motor vehicles or road dust is consistently associated with increases in the likelihood of respiratory symptoms or inhaler use (Table 5). Reduced likelihood of wheeze or inhaler use is associated with 5- $\mu\text{g}/\text{m}^3$ increases in the same day, but not for lagged exposures to the sulfur source. Significant reductions in the likelihood of wheeze were also associated with exposures to the biomass burning source.

The effect of adding a gaseous co-pollutant (NO₂, CO, SO₂, or O₃) to the source exposure




Figure 1. Daily prevalence of wheeze (A) and short-acting inhaler use (B). Daily PM_{2.5} source concentrations of motor vehicle (C), road dust (D), sulfur (E), biomass burning (F), oil (G), and sea salt (H). Dotted line at left of x-axis indicates warmer months (April–September). $n = 149$ children, New Haven, Connecticut, August 2000–January 2004.

model is shown for wheeze and short-acting inhaler use in Table 6. For example, the addition of any of the gaseous co-pollutants to the source exposure model reduces the effect of the motor vehicle source on wheeze. Interestingly, in the source model that also includes NO₂ (another marker for traffic), each 20-ppb increase in NO₂ increases the likelihood of wheeze by 8% ($p = 0.09$). No significant associations were seen in co-pollutant models between exposure and likelihood of persistent cough or chest tightness.

Discussion

Results of the source apportionment analysis suggest that over half of the mean PM_{2.5} can be attributed to traffic sources motor vehicles and road dust, and that daily exposure to these sources is associated with increased risk of respiratory symptoms and inhaler use in children with asthma. Risks remain in models that include all six PM_{2.5} sources as well as one gaseous co-pollutant. Of co-pollutants added to the model one at a time, NO₂, which is a marker for traffic (Grieshop et al. 2006), was also found to be an independent risk factor for increased wheeze (Table 6).

In single-element analyses, EC, also considered a marker for traffic (Grieshop et al. 2006; Sarnat et al. 2008), is itself associated with increased symptoms (wheeze, shortness of breath, chest tightness) and medication use (Table 4). Of the EC measured in PM_{2.5}, 88% was contributed by the traffic-related sources: 62% from motor vehicles and 26% from road dust (Table 3). Among elements associated with the road dust source, Ca appears to have the strongest association with symptoms (Table 4). Numerous studies have

found that the composition of Ca in road dust is higher than that in the earth's crust (Grieshop et al. 2006). Results from tunnel and chassis dynamometer studies have shown that Ca is emitted from light and heavy-duty vehicles and is possibly associated with the combustion of motor oil (Ning et al. 2008). The stronger associations with fine-particle Ca compared with other major terrestrial elements such as Al and Si suggests that it may be a better tracer of urban road dust.

A negative association was observed between biomass burning and wheeze (Table 5). Potassium, the tracer element for biomass burning source and an element used in firework propellant mixture, was present at high concentrations when Fourth of July displays took place (Moreno et al. 2007). When the analysis was rerun without the days with the five highest potassium values [all > 1,000 ng/m³ and all occurring in July (4 July 2001, 2002, 2003; 5 July 2003; 7 July 2002) (Figure 1F)], the negative association seen previously for wheeze remains but is no longer significant [odds ratio (OR) 0.85; $p = 0.29$]. Of note is the high level of potassium measured on 7 July 2002, which is attributable not to fireworks but to a smoke plume from a massive Canadian forest fire (Sigler et al. 2003). For our subjects, symptoms and medication use are at their lowest in summer and highest in winter (Figure 1A,B), and if these children with asthma were kept away from fireworks displays and/or kept indoors during the smoke plume days, this may have resulted in the apparent protective association with potassium levels.

Subjects who were in follow-up during the summer of 2001 also contributed data to

an earlier analysis of O₃ and respiratory health in southern New England (Gent et al. 2003). In that analysis, we found O₃ exposure to be a risk for increased respiratory symptoms, but only among children taking asthma maintenance medication (e.g., corticosteroids). The model with PM_{2.5} sources and O₃ shown in Table 6 shows O₃ to be a risk factor increasing the likelihood of wheeze by 8% ($p = 0.08$), but this model includes data from all children living in New Haven regardless of maintenance medication status. Reanalysis stratified by maintenance medication use showed that every 50-ppb increase in O₃ (8-hr average) resulted in a larger risk of wheeze for users of maintenance medication ($n = 76$) [OR = 1.29; 95% confidence interval (CI), 0.99–1.68; $p = 0.06$] than for nonusers ($n = 63$) (OR = 1.10; 95% CI, 0.77–1.57; $p = 0.60$). Comparable results from the earlier report are given for a 50-ppb increase in 1-hr O₃: For children on maintenance medication ($n = 130$), the likelihood of wheeze increases by 35% (OR = 1.35; 95% CI, 1.11–1.65) (Gent et al. 2003).

Although exposure to total PM_{2.5} mass has itself been shown to be a significant risk factor for adverse respiratory outcomes among children (Johnston et al. 2006; Mar et al. 2004; Rabinovitch et al. 2006; Ranzi et al. 2004; Rodriguez et al. 2007; Slaughter et al. 2003), this is not always the case (Chimonas and Gessner 2007; Holguin et al. 2007). In a study of Medicaid claims for doctor visits for asthma in Anchorage, Alaska, the authors noted that the major constituent of their particles is geologic in origin (i.e., rock dust), and that annual levels of PM mass are low, 6 µg/m³ (95% CI, 0.5–7.0). The authors found no association between exposure to PM_{2.5} mass

Table 3. Source contributions^a to PM_{2.5} mass (µg/m³) and elemental (ng/m³) concentrations in New Haven, Connecticut, August 2000 – January 2004.

Source	Source						Estimated	Mass	%RMSE
	Motor vehicles	Road dust	Sulfur	Biomass burning	Oil	Sea salt		Measured	
Element (ng/m ³)									
EC	961.3 ^b	403.4	118.8	22.1	68.1	−19.0	1554.8	1894.5	17.9
Zn	22.3 ^b	2.6	0.8	0.5	1.7	1.0	29.0	23.0	26.3
Pb	3.4 ^b	0.5	0.6	0.5	—	0.1	5.1	4.9	4.0
Cu	2.1 ^b	1.3	0.5	1.8	0.2	—	6.0	6.3	5.5
Se	0.4 ^b	−0.1	0.2	—	0.0	0.0	0.5	0.5	3.5
Si	−4.5	115.7 ^b	5.7	4.8	2.0	2.2	125.9	117.0	7.6
Fe	84.7	101.1 ^b	3.0	11.6	12.2	−1.7	210.9	219.0	3.7
Al	−7.0	54.8 ^b	7.0	5.8	—	1.6	62.0	67.5	8.1
Ca	17.8	29.1 ^b	—	1.5	1.1	2.0	51.6	51.2	0.7
Ba	2.9	4.9 ^b	0.5	4.1	0.8	−0.4	12.7	12.6	0.4
Ti	1.2	4.2 ^b	0.5	0.9	—	—	6.9	7.4	6.3
S	210.4	166.3	906.7 ^b	75.4	18.7	26.0	1403.4	1451.8	3.3
P	14.0	6.6	36.8 ^b	2.6	1.5	—	61.5	64.8	5.2
K	−8.0	13.8	12.4	60.5 ^b	−2.7	3.7	79.6	63.2	25.8
V	0.6	1.1	0.8	0.2	6.0 ^b	0.2	8.9	9.9	9.4
Ni	1.9	0.4	0.2	0.1	2.6 ^b	0.2	5.6	5.2	6.9
Na	20.9	29.3	37.0	4.4	—	47.0 ^b	140.1	180.8	22.5
Cl	20.5	—	−11.2	—	6.9	32.0 ^b	48.7	29.6	64.4
PM _{2.5} (µg/m ³)	6.6 ^b	2.3	5.5	0.9	0.8	0.5	16.6	17.0	2.4
Daily source contribution to PM _{2.5} (%)									
Median	42.5	12.3	28.7	4.4	2.2	1.2			
IQR	26.7	13.8	30.3	5.6	7.7	7.2			

—, nonsignificant contribution.

^aMean concentration for 1,181 days between August 2000 and January 2004. ^bMaximum contribution from a particular source for each element.

and health outcomes. On the other hand, significant effects were seen between increases in PM₁₀ mass concentrations (PM with aerodynamic diameter < 10 µm) and increased visits for asthma and prescriptions for inhalers (Chimonas and Gessner 2007). In a study of asthmatic children and respiratory health in a busy, traffic-filled Mexican border town, Holguin et al. (2007) found no significant association between level of PM_{2.5} mass and respiratory health. However, using traffic as the exposure metric (calculated as traffic density in 50-m residential buffers) revealed a significant risk for increased respiratory symptoms with increasing traffic (OR = 1.58; 95% CI, 1.05–2.38) (Holguin et al. 2007).

Studies using proximity to traffic as an exposure metric suggest that children

living near busy roads have impaired respiratory health. Large studies of schoolchildren in Munich (Nicolai et al. 2003) and Nottingham, United Kingdom (Venn et al. 2001), have found that proximity of residence (< 50–90 m) to heavily traveled roads is associated with increased risk of asthma symptoms. Studies of schoolchildren conducted in the Netherlands found increased health risks associated with proximity specifically to truck traffic and wheeze (Janssen et al. 2003), chronic respiratory symptoms (Van Vliet et al. 1997), and lung function (Brunekreef et al. 1997). Similar increased risk associations were found between black smoke (a proxy for diesel exhaust), measured at subjects' schools, and respiratory outcomes. No significant associations were found between

respiratory symptoms and volume of automobile traffic. The strongest associations were found for children living within 300 m of a highway (Brunekreef et al. 1997). A smaller study in Cincinnati, Ohio (Ryan et al. 2005), found an increased risk of wheezing in infants associated with stop-and-go bus and truck traffic within 100 m of the residence. All of these studies suggest that proximity to traffic, in particular high-volume truck traffic, has a negative impact on the respiratory health of children. Rather than a metric based on proximity to roads or traffic density, we used factor analysis and source apportionment techniques to examine exposures to potentially toxic portions of particle mass and the health of children residing in an area of significant mobile source activity.

Table 4. ORs from separate repeated measures logistic regression analyses^a of associations between daily respiratory symptoms and each daily elemental concentration of PM_{2.5}.

Source/element	Lag	Unit increase (ng/m ³)	Wheeze		Persistent cough		Shortness of breath		Chest tightness		Inhaler short-acting	
			OR	p-Value	OR	p-Value	OR	p-Value	OR	p-Value	OR	p-Value
Motor vehicles												
EC	L0	1,000	1.04	0.04	1.01	0.42	1.06	0.001	1.03	0.20	1.01	0.15
	L02	1,000	1.07	0.06	1.03	0.23	1.12	0.01	1.10	0.04	1.02	0.40
Zn	L0	10	1.00	0.69	1.00	0.60	1.02	0.001	1.00	0.72	1.00	0.41
	L02	10	1.00	0.98	1.00	0.94	1.04	0.06	1.03	0.13	1.01	0.53
Pb	L0	5	1.02	0.31	1.02	0.25	1.03	0.11	1.02	0.31	1.01	0.06
	L02	5	1.07	0.13	1.05	0.12	1.12	0.01	1.10	0.02	1.04	0.10
Cu	L0	5	1.01	0.59	1.02	0.13	1.06	0.01	1.03	0.23	1.01	0.22
	L02	5	1.02	0.67	1.05	0.04	1.06	0.21	1.04	0.39	1.01	0.46
Se	L0	1	1.00	0.97	1.00	0.84	1.02	0.40	1.00	0.79	0.99	0.20
	L02	1	1.02	0.71	0.98	0.43	1.02	0.67	0.98	0.61	0.99	0.75
Road dust												
Si	L0	100	1.03	0.03	1.02	0.01	1.04	0.01	1.02	0.20	1.02	0.004
	L02	100	1.07	0.04	1.05	0.02	1.08	0.02	1.06	0.10	1.03	0.09
Fe	L0	100	1.04	0.02	1.02	0.06	1.06	0.002	1.01	0.47	1.02	0.004
	L02	100	1.07	0.05	1.04	0.04	1.08	0.04	1.05	0.21	1.03	0.08
Al	L0	50	1.02	0.17	1.03	0.002	1.05	0.002	1.02	0.21	1.02	0.02
	L02	50	1.07	0.03	1.06	0.01	1.09	0.004	1.07	0.04	1.02	0.11
Ca	L0	50	1.07	0.02	1.05	0.01	1.10	0.002	1.04	0.26	1.04	0.01
	L02	50	1.14	0.04	1.09	0.03	1.18	0.01	1.14	0.07	1.04	0.17
Ba	L0	10	0.99	0.57	1.00	0.83	1.04	0.02	1.01	0.63	1.01	0.08
	L02	10	0.99	0.81	1.00	0.81	1.03	0.38	1.02	0.51	1.01	0.36
Ti	L0	5	1.00	0.59	1.00	0.57	1.01	0.01	1.00	0.34	1.00	0.72
	L02	5	1.01	0.56	1.01	0.29	1.03	0.05	1.01	0.52	1.00	0.66
Sulfur												
S	L0	1,000	0.98	0.43	1.00	0.84	1.01	0.63	0.99	0.80	0.99	0.13
	L02	1,000	1.00	0.99	1.02	0.27	1.01	0.79	1.02	0.68	1.00	0.81
P	L0	50	0.98	0.39	1.00	0.75	1.01	0.61	1.00	0.88	0.98	0.15
	L02	50	0.99	0.89	1.03	0.30	1.01	0.78	1.02	0.67	1.00	0.99
Biomass burning												
K	L0	50	0.98	0.06	1.00	0.64	1.01	0.01	1.01	0.02	1.00	0.68
	L02	50	0.96	0.04	1.00	0.86	1.00	0.79	0.99	0.67	0.99	0.28
Oil												
V	L0	10	0.99	0.73	1.01	0.56	1.01	0.46	0.99	0.71	0.98	0.12
	L02	10	0.93	0.04	0.96	0.05	0.98	0.58	0.94	0.12	0.96	0.03
Ni	L0	5	1.01	0.59	1.01	0.21	1.04	0.05	1.01	0.58	1.01	0.48
	L02	5	0.99	0.72	1.00	0.99	1.04	0.32	1.01	0.84	1.01	0.48
Sea salt												
Na	L0	100	0.98	0.23	1.00	0.58	1.00	0.94	0.99	0.43	0.99	0.35
	L02	100	0.97	0.29	0.98	0.21	0.99	0.74	0.98	0.61	0.99	0.37
Cl	L0	10	1.00	0.89	1.00	0.31	1.00	0.89	1.00	0.24	1.00	0.69
	L02	10	1.00	0.81	1.00	0.06	1.00	0.80	1.00	0.65	1.00	0.83

Associations are shown for same-day exposure (L0) and averaged over the same day and previous 2 days (L02). *n* = 149 children with asthma, New Haven, Connecticut, August 2000–January 2004.

^aSeparate analyses were performed for each element and each respiratory symptom and medication use. Models were adjusted for season, day of week, and date. ORs are given for the unit increase given in the second column for each element (and EC). Individual elements in fine particle mass may come from more than one source but are listed here according to their major source. See Table 3 for the contribution of each source to each element's total mass. Results from analyses using elemental exposures lagged by 1 or 2 days (L1, L2) are included in Supplemental Material, Table 2 (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>).

One strength of this study is its longitudinal design. More than 46,000 subject-days of data were available for analyses (resulting in small CIs) where each subject served as his or her own control for potential confounders. Effect estimates from analyses where sex, age, ethnicity, parental education level, and exposure to environmental tobacco smoke were also included varied little from those shown in Table 6. For example, each 5- $\mu\text{g}/\text{m}^3$ increase in exposure to motor vehicle source was associated with identical increased risks of wheeze and shortness of breath and inhaler use. Likewise, the risk estimates associated with exposure to road dust source were identical for persistent cough and inhaler use and slightly decreased for wheeze (OR = 1.09; 95% CI, 1.01–1.19) and shortness of breath (OR = 1.11; 95% CI,

1.02–1.21) (for comparison, see Table 5). Goodness-of-fit tests suggest that the models for wheeze and chest tightness are reasonably good fits to the data (Table 5). There were no systematic patterns to the lack of fit for models for persistent cough, shortness of breath, or short-acting inhaler use. However, because of the repeated measurements, observations were not independent in any of the models, which may affect the interpretation of the Hosmer–Lemeshow statistic. It is possible that the more frequently reported events of persistent cough and short-acting inhaler use (Table 2) may be associated with ambient air pollution in combination with other factors (e.g., activity level) not included in the current study.

One limitation of our study is the lack of co-located EC calibration measurements.

Previous studies have used thermal optical reflectance (TOR) to measure EC collected on quartz filters in co-located monitors. TOR, a relatively costly but accurate measure of EC, is used to calibrate EC estimated by the much less expensive light reflectance measures on Teflon filters. We assumed a factor of 1 based on calibration factors reported in other studies examining air pollution in urban settings (Cyrus et al. 2003; Kinney et al. 2000) and on our own results of TOR EC measurements on 30, co-located quartz filters [see Supplemental Material (available online at <http://www.ehponline.org/members/2009/0800335/suppl.pdf>)].

Another limitation to our study is our reliance on a central site monitor. It is likely that measurements from this site, located on an on-ramp to I-95 at the southern edge of New Haven County, represent the highest end of the exposure distribution. Exposure error for regional sources, for example, sulfur, is probably low because of its homogeneity within the study area. On the other hand, local traffic exposures vary across the study area, which may have led to an overestimate of subjects' exposures to motor vehicle and road dust sources and an underestimate of health effects; that is, it is possible that levels lower than those used have significant adverse effects on respiratory health of children with asthma in our region. Determination of spatial variability of $\text{PM}_{2.5}$ away from central site monitors will be needed to improve assessment of exposures actually experienced by vulnerable populations.

Conclusions

The composition of ambient fine particles is complex, because it depends on emission sources and atmospheric processes. The U.S. EPA national air quality standard for $\text{PM}_{2.5}$ considers only mass and not composition (U.S. EPA 2009). However, health impacts associated with exposures to specific particle components or sources, especially for vulnerable populations, may be missed with this aggregate characterization (Franklin et al. 2008; Sarnat et al. 2008). Apportioning fine particle mass according to contributing sources may help unmask source-specific associations. For children with asthma living in an area of noncompliance with $\text{PM}_{2.5}$ standards, increased risk of daily symptoms and medication use was associated with daily traffic-related fine particle sources.

REFERENCES

- Andersen ZJ, Wahlin P, Raaschou-Nielsen O, Scheike T, Loft S. 2007. Ambient particle source apportionment and daily hospital admissions among children and elderly in Copenhagen. *J Exp Sci Environ Epidemiol* 17(7):625–636.
- Belanger K, Beckett W, Triche E, Bracken MB, Holford T, Ren P, et al. 2003. Symptoms of wheeze and persistent cough in the first year of life: associations with indoor allergens, air contaminants and maternal history of asthma. *Am J Epidemiol* 158:195–202.

Table 5. ORs (95% CIs) from repeated measures logistic regression models of respiratory symptoms and rescue medication use and daily source concentrations of $\text{PM}_{2.5}$.

Source	Wheeze	Persistent cough	Shortness of breath	Chest tightness	Inhaler short-acting
L0 model					
Motor vehicles	1.05 (0.99–1.10)	1.02 (0.99–1.04)	1.06 (1.01–1.11)	1.02 (0.97–1.08)	1.02 (1.00–1.05)
Road dust	1.10 (1.01–1.19)	1.06 (1.01–1.11)	1.12 (1.02–1.22)	1.04 (0.95–1.15)	1.06 (1.02–1.11)
Sulfur	0.97 (0.94–1.00)	1.00 (0.98–1.01)	0.98 (0.94–1.02)	0.99 (0.94–1.03)	0.98 (0.97–1.00)
Biomass burning	0.80 (0.66–0.98)	0.97 (0.92–1.03)	1.05 (0.95–1.17)	1.06 (0.95–1.18)	1.00 (0.96–1.03)
Oil	1.02 (0.86–1.20)	1.02 (0.95–1.10)	1.07 (0.92–1.26)	0.99 (0.82–1.18)	0.98 (0.91–1.05)
Sea salt	0.96 (0.86–1.07)	0.99 (0.92–1.07)	1.01 (0.92–1.12)	0.95 (0.84–1.08)	0.99 (0.94–1.04)
L02 model					
Motor vehicles	1.10 (1.01–1.19)	1.03 (0.98–1.09)	1.12 (1.01–1.24)	1.08 (0.98–1.20)	1.03 (0.98–1.08)
Road dust	1.26 (1.05–1.51)	1.16 (1.02–1.32)	1.28 (1.05–1.55)	1.20 (0.97–1.49)	1.09 (1.00–1.19)
Sulfur	0.98 (0.92–1.04)	1.01 (0.98–1.05)	0.97 (0.90–1.04)	1.00 (0.92–1.08)	1.00 (0.97–1.03)
Biomass burning	0.64 (0.46–0.88)	0.93 (0.81–1.06)	0.78 (0.52–1.18)	0.87 (0.62–1.22)	0.95 (0.87–1.04)
Oil	0.80 (0.56–1.08)	0.84 (0.71–1.00)	0.94 (0.69–1.29)	0.80 (0.58–1.10)	0.92 (0.81–1.05)
Sea salt	0.91 (0.82–1.16)	0.88 (0.77–1.01)	1.01 (0.79–1.29)	0.95 (0.71–1.27)	0.97 (0.88–1.07)

$n = 149$ children with asthma, New Haven, Connecticut, August 2000–January 2004. Separate logistic regression analyses were performed for each health outcome. Each model included the six $\text{PM}_{2.5}$ sources listed, either as same-day concentrations (L0), or the concentration averaged over the same day and previous 2 days (L02), as well as variables controlling for season, day of week, and date. Logistic regressions were performed using GEE. All ORs are given for a 5- $\mu\text{g}/\text{m}^3$ increase in source concentration.

Table 6. Effect of adding a gaseous co-pollutant to the source exposure model [OR (95% CI)].

Outcome/source	Sources alone	Co-pollutants			
		NO ₂	CO	SO ₂	O ₃
Wheeze					
Motor vehicles	1.05 (0.99–1.10)	1.03 (0.98–1.08)	1.05 (0.99–1.11)	1.04 (0.99–1.09)	1.06 (0.97–1.16)
Road dust	1.10 (1.01–1.19)	1.11 (1.02–1.20)	1.10 (1.01–1.19)	1.10 (1.01–1.19)	1.11 (1.01–1.23)
Sulfur	0.97 (0.94–1.00)	0.96 (0.92–0.99)	0.97 (0.94–1.01)	0.97 (0.93–1.00)	0.95 (0.91–1.00)
Biomass burning	0.80 (0.66–0.98)	0.79 (0.65–0.98)	0.80 (0.66–0.98)	0.79 (0.64–0.98)	0.74 (0.57–0.97)
Oil	1.02 (0.86–1.20)	1.02 (0.87–1.21)	1.02 (0.86–1.20)	1.01 (0.86–1.19)	0.92 (0.62–1.39)
Sea salt	0.96 (0.86–1.07)	0.96 (0.85–1.07)	0.96 (0.86–1.08)	0.95 (0.85–1.07)	1.01 (0.72–1.40)
Co-pollutant		1.08 (0.99–1.18)	1.00 (0.94–1.07)	1.02 (0.96–1.09)	1.08 (0.99–1.18)
Short-acting inhaler use					
Motor vehicles	1.02 (1.00–1.05)	1.02 (0.99–1.04)	1.02 (0.99–1.05)	1.02 (0.99–1.04)	1.02 (0.98–1.07)
Road dust	1.06 (1.02–1.11)	1.06 (1.02–1.10)	1.06 (1.02–1.11)	1.06 (1.02–1.11)	1.06 (1.00–1.13)
Sulfur	0.98 (0.97–1.00)	0.98 (0.96–1.00)	0.98 (0.96–1.00)	0.98 (0.96–1.00)	0.97 (0.95–1.00)
Biomass burning	1.00 (0.96–1.03)	1.00 (0.96–1.03)	0.99 (0.96–1.03)	0.99 (0.96–1.03)	0.99 (0.95–1.03)
Oil	0.98 (0.91–1.05)	0.98 (0.91–1.05)	0.97 (0.91–1.04)	0.97 (0.91–1.04)	1.03 (0.88–1.22)
Sea salt	0.99 (0.94–1.04)	0.99 (0.94–1.04)	0.99 (0.94–1.04)	0.99 (0.94–1.04)	1.01 (0.88–1.15)
Co-pollutant		1.01 (0.97–1.06)	1.02 (0.98–1.05)	1.01 (0.99–1.04)	1.01 (0.97–1.05)

$n = 149$ children with asthma, New Haven, Connecticut, August 2000–January 2004. Separate repeated measures logistic regression analyses using GEE were performed for each health outcome. Each model includes the six particle sources (same-day exposure) as well as season, day of week, and date. Gaseous co-pollutants were added to the source exposure model one at a time. ORs for sources are given for a 5- $\mu\text{g}/\text{m}^3$ increase in source concentration; ORs for co-pollutants are given for each 20 ppb (NO_2 , ozone), 0.5 ppm (CO), or 5 ppb (SO_2). Pearson correlations between $\text{PM}_{2.5}$ sources and criteria pollutants were < 0.20 except for motor vehicles with NO_2 ($r = 0.49$), CO ($r = 0.59$), SO_2 ($r = 0.45$); road dust with NO_2 ($r = 0.37$), O_3 ($r = 0.58$); and oil with CO ($r = 0.22$), SO_2 ($r = 0.32$).

- Brown SG, Frankel A, Raffuse SM, Roberts PT, Hafner HR, Anderson DJ. 2007. Source apportionment of fine particulate matter in Phoenix, AZ, using positive matrix factorization. *J Air Waste Manag Assoc* 57:741–752.
- Brunekreef B, Janssen NAH, de Hartog JJ, Harssema H, Knafe M, Van Vliet P. 1997. Air pollution from truck traffic and lung function in children living near motorways. *Epidemiology* 8:298–303.
- Chardon B, Lefranc A, Granados D, Gremy I. 2007. Air pollution and doctors' house calls for respiratory diseases in the Greater Paris area (2000–2003). *Occup Environ Med* 64:320–324.
- Chimonas M-AR, Gessner BD. 2007. Airborne particulate matter from primarily geologic, non-industrial sources at levels below National Ambient Air Quality Standards is associated with outpatient visits for asthma and quick-relief medication prescriptions among children less than 20 years old enrolled in Medicaid in Anchorage, Alaska. *Environ Research* 103:397–404.
- Cyrys J, Heinrich J, Hoek G, Meliefste K, Lewne M, Gehring U, et al. 2003. Comparison between different traffic-related particle indicators: elemental carbon (EC), PM_{2.5} mass, and absorbance. *J Exp Anal Environ Epidemiol* 13:134–143.
- Delfino RJ, Quintana PJE, Floro J, Gastanaga VM, Samimi BS, Kleinman MT, et al. 2004. Association of FEV₁ in asthmatic children with personal and microenvironmental exposure to airborne particulate matter. *Environ Health Perspect* 112:932–941.
- Franklin M, Koutrakis P, Schwartz J. 2008. The role of particle composition on the association between PM_{2.5} and mortality. *Epidemiology* 19:680–689.
- Gent JF, Koutrakis P, Leaderer BP. 2008. Childhood asthma: association of exposure to fine particle sources with respiratory symptoms [Abstract]. *Am J Respir Crit Care Med* 177:A186.
- Gent JF, Triche EW, Holford TR, Belanger K, Bracken MB, Beckett WS, et al. 2003. Association of low-level ozone and fine particles with respiratory symptoms in children with asthma. *JAMA* 290:1859–1867.
- Grieshop AP, Lipsky EM, Pekney NJ, Takahama S, Robinson AL. 2006. Fine particle emission factors from vehicles in a highway tunnel: effects of fleet composition and season. *Atmos Environ* 40:S287–S298.
- Hertz-Picciotto I, Baker RJ, Yap P-S, Dostal M, Joad JP, Lipsett M, et al. 2007. Early childhood lower respiratory illness and air pollution. *Environ Health Perspect* 115:1510–1518.
- Holguin F, Flores S, Ross Z, Corez M, Molina M, Molina L, et al. 2007. Traffic-related exposures, airway function, inflammation, and respiratory symptoms in children. *Am J Respir Crit Care Med* 176(12):1236–1242.
- Hong Y-C, Hwang S-S, Kim JH, Lee K-H, Lee H-J, Lee K-H, et al. 2007. Metals in particulate pollutants affect peak expiratory flow of schoolchildren. *Environ Health Perspect* 115:430–434.
- Janssen NAH, Brunekreef B, Van Vliet P, Aarts F, Meliefste K, Harssema H, et al. 2003. The relationship between air pollution from heavy traffic and allergic sensitization, bronchial hyperresponsiveness, and respiratory symptoms in Dutch schoolchildren. *Environ Health Perspect* 111:1512–1518.
- Janssen NAH, van Vliet PHN, Aarts F, Harssema H, Brunekreef B. 2001. Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmos Environ* 35:3875–3884.
- Johnston FH, Webby RJ, Pilotto LS, Bailie RS, Parry DL, Halpin SJ. 2006. Vegetation fires, particulate air pollution and asthma: a panel study in the Australian monsoon tropics. *Int J Environ Health Res* 16(6):391–404.
- Kavouras IG, Koutrakis P, Cereceda-Balic F, Oyola P. 2001. Source apportionment of PM₁₀ and PM_{2.5} in five Chilean cities using factor analysis. *J Air Waste Manag Assoc* 51:451–464.
- Kinney PL, Aggarwal M, Northridge ME, Hanssen NAH, Shepard P. 2000. Airborne concentrations of PM_{2.5} and diesel exhaust particles on Harlem sidewalks: a community-based pilot study. *Environ Health Perspect* 108:213–218.
- Laden F, Neas LM, Dockery DW, Schwartz J. 2000. Association of fine particulate matter from different sources with daily mortality in six U.S. cities. *Environ Health Perspect* 108:941–947.
- Lewis TC, Robins TG, Dvorchak JT, Keeler GJ, Yip FY, Mentz GB, et al. 2005. Air pollution-associated changes in lung function among asthmatic children in Detroit. *Environ Health Perspect* 113:1068–1075.
- Mar TF, Larson TV, Stier RA, Claiborn C, Koenig JQ. 2004. An analysis of the association between respiratory symptoms in subjects with asthma and daily air pollution in Spokane, Washington. *Inhal Toxicol* 16:809–815.
- Mar TF, Norris GA, Koenig JQ, Larson TV. 2000. Associations between air pollution and mortality in Phoenix, 1995–1997. *Environ Health Perspect* 108:347–353.
- Moreno T, Querol X, Alastuey A, Minguillon M, Pey J, Rodriguez S, et al. 2007. Recreational atmospheric pollution episodes: Inhalable metalliferous particles from firework displays. *Atmos Environ* 41:913–922.
- Moshhammer H, Hutter H-P, Hauck H, Neuberger M. 2006. Low levels of air pollution induce changes of lung function in a panel of schoolchildren. *Eur Respir J* 27(6):1138–1143.
- Nicolai T, Carr D, Weiland SK, Duhme H, von Ehrenstein O, Wagner C, et al. 2003. Urban traffic and pollutant exposure related to respiratory outcomes and atopy in a large sample of children. *Eur Respir J* 21:956–963.
- Ning Z, Polidori A, Schauer JJ, Sioutas C. 2008. Emission factors of PM species based on freeway measurements and comparison with tunnel and dynamometer studies. *Atmos Environ* 42:3099–3114.
- Ozkaynak H, Thurston GD. 1987. Associations between 1980 U.S. mortality rates and alternative measures of airborne particle concentration. *Risk Anal* 7(4):449–461.
- Penttinen P, Vallius M, Tiittanen P, Ruuskanen J, Pekkanen J. 2006. Source-specific fine particles in urban air and respiratory function among adult asthmatics. *Inhal Toxicol* 18:191–198.
- Pino P, Walter T, Oyarzun M, Villegas R, Romieu I. 2004. Fine particulate matter and wheezing illnesses in the first year of life. *Epidemiology* 15(6):702–708.
- Rabinovitch N, Strand M, Gelfand EW. 2006. Particulate levels are associated with early asthma worsening in children with persistent disease. *Am J Respir Crit Care Med* 173(10):1098–1105.
- Ranzi A, Gambini M, Spattini A, Galassi C, Sesti D, Bedeschi M, et al. 2004. Air pollution and respiratory status in asthmatic children: hints for a locally based preventive strategy. AIRE study. *Eur J Epidemiol* 19(6):567–576.
- Rodriguez C, Tonkin R, Heyworth J, Kusel M, DeKlerk N, Sly P, et al. 2007. The relationship between outdoor air quality and respiratory symptoms in young children. *Int J Environ Health Res* 17(5):351–360.
- Ryan PH, LeMasters G, Biagini J, Bernstein D, Grinshpun SA, Shukla R, et al. 2005. Is it traffic type, volume, or distance? Wheezing in infants living near truck and bus traffic. *J Allerg Clin Immunol* 116:279–284.
- Sarnat JA, Marmur A, Klein M, Kim E, Russell AG, Sarnat SE, et al. 2008. Fine particle sources and cardiorespiratory morbidity: an application of chemical mass balance and factor analytical source-apportionment methods. *Environ Health Perspect* 116:459–466.
- Sigler JM, Lee X, Munger W. 2003. Emission and long-range transport of gaseous mercury from a large-scale Canadian boreal forest fire. *Environ Sci Tech* 37(19):4343–4347.
- Slaughter JC, Lumley T, Sheppard L, Koenig JQ, Shapiro GG. 2003. Effects of ambient air pollution on symptom severity and medication use in children with asthma. *Ann Allergy Asthma Immunol* 91(4):346–353.
- Taino M, Tuomisto JT, Hanninen O, Aarnio P, Koistinen KJ, Jantunen MJ, et al. 2005. Health effects caused by primary fine particulate matter (PM_{2.5}) emitted from buses in the Helsinki metropolitan area, Finland. *Risk Anal* 25(1):151–160.
- Thurston GD, Spengler JD. 1985. A quantitative assessment of source contributions to inhalable particulate matter pollution in metropolitan Boston. *Atmos Environ* 19(1):9–25.
- Trenga CA, Sullivan JH, Schildcrout JS, Shepherd KP, Shapiro GG, Liu LJS, et al. 2006. Effect of particulate air pollution on lung function in adult and pediatric subjects in a Seattle panel study. *Chest* 129(6):1614–1622.
- U.S. Census Bureau. 2000. United States Census 2000. Available: <http://www.census.gov/main/www/cen2000.html> [accessed 1 April 2008].
- U.S. Department of Health and Human Services. 2002. Global Initiative for Asthma, Global Strategy for Asthma Management and Prevention 02-3659. Washington, DC:National Institutes of Health, National Heart, Lung, and Blood Institute.
- U.S. EPA (U.S. Environmental Protection Agency). 2008. National Emissions Inventory (NEI). Available: <http://www.epa.gov/ttn/chieff/sofware/speciate/index.html> [accessed 1 June 2009].
- U.S. EPA (U.S. Environmental Protection Agency). 2009. National Ambient Air Quality Standards (NAAQS) for Particulate Matter (PM). Available: http://www.epa.gov/ttn/naaqs/standards/pm/s_pm_index.html [accessed 1 January 2009].
- Van Vliet P, Knafe M, de Hartog J, Janssen N, Harssema H, Brunekreef B. 1997. Motor vehicle exhaust and chronic respiratory symptoms in children living near freeways. *Environ Res* 74:122–132.
- Venn AJ, Lewis SA, Cooper M, Hubbard R, Britton J. 2001. Living near a main road and the risk of wheezing illness in children. *Am J Respir Crit Care Med* 164:2177–2180.
- Watson JG, Chow JC, Frazier CA. 1999. X-ray fluorescence analysis of ambient air samples. In: *Elemental Analysis of Airborne Particles* (Landsberger S, Creatchman M, eds). Amsterdam: Gordon and Breach Science Publishers, 67–96.
- Watson JG, Zhu T, Chow JC, Engelbrecht J, Fujita EM, Wilson WE. 2002. Receptor modeling application framework for particle source apportionment. *Chemosphere* 49:1093–1136.